

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup>:</b> <b>C07K 14/47, C12P 21/02</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/27878</b> <b>(43) International Publication Date:</b> 18 May 2000 (18.05.00)
<b>(21) International Application Number:</b> PCT/GB99/03730 <b>(22) International Filing Date:</b> 9 November 1999 (09.11.99)  <b>(30) Priority Data:</b> 9824544.2      9 November 1998 (09.11.98)      GB  <b>(71) Applicant (for all designated States except US):</b> GENDAQ LIMITED [GB/GB]; 1-3 Burtonhole Lane, Mill Hill, London NW7 1AD (GB).  <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> CHOO, Yen [GR/GB]; MRC Laboratory of Molecular Biology, Medical Research Council Centre, Hills Road, Cambridge CB2 2QH (GB).  <b>(74) Agents:</b> MASCHIO, Antonio et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> SCREENING SYSTEM FOR ZINC FINGER POLYPEPTIDES FOR A DESIRED BINDING ABILITY  <b>(57) Abstract</b>  The invention relates to a method for producing a zinc finger nucleic acid binding protein comprising preparing a zinc finger protein according design rules, varying the protein at one or more positions, and selecting variants which bind to a target nucleic acid sequence by polysome display.		

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## SCREENING SYSTEM FOR ZINC FINGER POLYPEPTIDES FOR A DESIRED BINDING ABILITY

The present application relates to a method for screening zinc finger polypeptides for a desired binding ability. In particular, the invention relates to a polysome display  
5 technique which permits the isolation of binding polypeptides without resorting to phage display techniques.

Protein-nucleic acid recognition is a commonplace phenomenon which is central to a large number of biomolecular control mechanisms which regulate the functioning of  
10 eukaryotic and prokaryotic cells. For instance, protein-DNA interactions form the basis of the regulation of gene expression and are thus one of the subjects most widely studied by molecular biologists.

A wealth of biochemical and structural information explains the details of protein-DNA  
15 recognition in numerous instances, to the extent that general principles of recognition have emerged. Many DNA-binding proteins contain independently folded domains for the recognition of DNA, and these domains in turn belong to a large number of structural families, such as the leucine zipper, the "helix-turn-helix" and zinc finger families.

20 Most sequence-specific DNA-binding proteins bind to the DNA double helix by inserting an  $\alpha$ -helix into the major groove (Pabo & Sauer 1992 *Annu. Rev. Biochem.* 61, 1053-1095; Harrison 1991 *Nature (London)* 353, 715-719; and Klug 1993 *Gene* 135, 83-92). Sequence specificity results from the geometrical and chemical complementarity  
25 between the amino acid side chains of the  $\alpha$ -helix and the accessible groups exposed on the edges of base-pairs. In addition to this direct reading of the DNA sequence, interactions with the DNA backbone stabilise the complex and are sensitive to the conformation of the nucleic acid, which in turn depends on the base sequence (Dickerson & Drew 1981 *J. Mol. Biol.* 149, 761-786. Crystal structures of protein-DNA complexes  
30 have shown that proteins can be idiosyncratic in their mode of DNA recognition, at least partly because they may use alternative geometries to present their sensory  $\alpha$ -helices to

DNA, allowing a variety of different base contacts to be made by a single amino acid and vice versa (Matthews 1988 Nature (London) 335, 294-295).

Protein engineering experiments have shown that it is possible to alter rationally the DNA-binding characteristics of individual zinc fingers when one or more of the  $\alpha$ -helical positions is varied in a number of proteins (Nardelli *et al.*, 1991 Nature (London) 349, 175-178; Nardelli *et al.*, 1992 Nucleic Acids Res. 20, 4137-4144; and Desjarlais & Berg 1992a Proteins 13, 272). It has already been possible to propose some principles relating amino acids on the  $\alpha$ -helix to corresponding bases in the bound DNA sequence (Desjarlais & Berg 1992b Proc. Natl. Acad. Sci. USA 89, 7345-7349). However in this approach the altered positions on the  $\alpha$ -helix are prejudged, making it possible to overlook the role of positions which are not currently considered important; and secondly, owing to the importance of context, concomitant alterations are sometimes required to affect specificity (Desjarlais & Berg 1992b), so that a significant correlation between an amino acid and base may be misconstrued.

More sophisticated principles describing the relationship between the sequence of the zinc finger and the nucleic acid target have been described, for example in WO 96/06166 (Medical Research Council).

20

To investigate binding of mutant Zf proteins, Thiesen and Bach (1991 FEBS 283, 23-26) mutated Zf fingers and studied their binding to randomised oligonucleotides, using electrophoretic mobility shift assays. Subsequent use of phage display technology has permitted the expression of random libraries of Zf mutant proteins on the surface of bacteriophage. The three Zf domains of Zif268, with 4 positions within finger one randomised, have been displayed on the surface of filamentous phage by Rebar and Pabo (1994 Science 263, 671-673). The library was then subjected to rounds of affinity selection by binding to target DNA oligonucleotide sequences in order to obtain Zf proteins with new binding specificities. Randomised mutagenesis (at the same positions as those selected by Rebar & Pabo) of finger 1 of Zif 268 with phage display has also been

30

used by Jamieson *et al.*, (1994 Biochemistry 33, 5689-5695) to create novel binding specificity and affinity.

In summary, it is known that Zf protein motifs are widespread in DNA binding proteins and that binding is via three key amino acids, each one contacting a single base pair in the target DNA sequence. Motifs are modular and may be linked together to form a set of fingers which recognise a contiguous DNA sequence (e.g. a three fingered protein will recognise a 9mer etc). The key residues involved in DNA binding have been identified through sequence data and from structural information. Directed and random mutagenesis has confirmed the role of these amino acids in determining specificity and affinity. Phage display has been used to screen for new binding specificities of random mutants of fingers. Therefore, the combination of a set of rules with a selection process appears to provide the most promising avenue for the development of zinc finger proteins.

#### 15 Summary of the Invention

According to a first aspect of the present invention, there is provided a method for producing a zinc finger nucleic acid binding protein comprising preparing a zinc finger protein according design rules, varying the protein at one or more positions, and selecting variants which bind to a target nucleic acid sequence by polysome display.

According to a second aspect of the present invention, there is provided a method for producing a zinc finger nucleic acid binding protein comprising an at least partially varied sequence and selecting variants thereof which bind to a target DNA strand, comprising the steps of:

- (i) preparing a nucleic acid binding protein of the Cys2-His2 zinc finger class capable of binding to a nucleic acid triplet in a target nucleic acid sequence, wherein binding to each base of the triplet by an  $\alpha$ -helical zinc finger nucleic acid binding motif in the protein is determined as follows:

- a) if the 5' base in the triplet is G, then position +6 in the  $\alpha$ -helix is Arg; or position +6 is Ser or Thr and position +2 is Asp;
- b) if the 5' base in the triplet is A, then position +6 in the  $\alpha$ -helix is Gln and +2 is not Asp;
- 5 c) if the 5' base in the triplet is T, then position +6 in the  $\alpha$ -helix is Ser or Thr and position +2 is Asp;
- d) if the 5' base in the triplet is C, then position +6 in the  $\alpha$ -helix may be any amino acid, provided that position +2 in the  $\alpha$ -helix is not Asp;
- e) if the central base in the triplet is G, then position +3 in the  $\alpha$ -helix is His;
- 10 f) if the central base in the triplet is A, then position +3 in the  $\alpha$ -helix is Asn;
- g) if the central base in the triplet is T, then position +3 in the  $\alpha$ -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;
- h) if the central base in the triplet is C, then position +3 in the  $\alpha$ -helix is Ser, Asp, Glu, Leu, Thr or Val;
- 15 i) if the 3' base in the triplet is G, then position -1 in the  $\alpha$ -helix is Arg;
- j) if the 3' base in the triplet is A, then position -1 in the  $\alpha$ -helix is Gln;
- k) if the 3' base in the triplet is T, then position -1 in the  $\alpha$ -helix is Asn or Gln;
- l) if the 3' base in the triplet is C, then position -1 in the  $\alpha$ -helix is Asp;
- 20 (ii) varying the resultant polypeptide at at least one position; and
- (iii) selecting the variants which bind to a target nucleic acid sequence by polysome display.

## 25 Detailed Description of the Invention

All of the nucleic acid-binding residue positions of zinc fingers, as referred to herein, are numbered from the first residue in the  $\alpha$ -helix of the finger, ranging from +1 to +9. "-1" refers to the residue in the framework structure immediately preceding the

30  $\alpha$ -helix in a Cys2-His2 zinc finger polypeptide. Residues referred to as "++" are

residues present in an adjacent (C-terminal) finger. Where there is no C-terminal adjacent finger, "++" interactions do not operate.

Cys2-His2 zinc finger binding proteins, as is well known in the art, bind to target  
5 nucleic acid sequences via  $\alpha$ -helical zinc metal atom co-ordinated binding motifs known  
as zinc fingers. Each zinc finger in a zinc finger nucleic acid binding protein is  
responsible for determining binding to a nucleic acid triplet in a nucleic acid binding  
sequence. Preferably, there are 2 or more zinc fingers, for example 2, 3, 4, 5 or 6 zinc  
fingers, in each binding protein. Advantageously, there are 3 zinc fingers in each zinc  
10 finger binding protein.

The method of the present invention allows the production of what are essentially  
artificial nucleic acid binding proteins. In these proteins, artificial analogues of amino  
acids may be used, to impart the proteins with desired properties or for other reasons.  
15 Thus, the term "amino acid", particularly in the context where "any amino acid" is  
referred to, means any sort of natural or artificial amino acid or amino acid analogue  
that may be employed in protein construction according to methods known in the art.  
Moreover, any specific amino acid referred to herein may be replaced by a functional  
analogue thereof, particularly an artificial functional analogue. The nomenclature used  
20 herein therefore specifically comprises within its scope functional analogues of the  
defined amino acids.

The  $\alpha$ -helix of a zinc finger binding protein aligns antiparallel to the nucleic acid  
strand, such that the primary nucleic acid sequence is arranged 3' to 5' in order to  
25 correspond with the N terminal to C-terminal sequence of the zinc finger. Since  
nucleic acid sequences are conventionally written 5' to 3', and amino acid sequences N-  
terminus to C-terminus, the result is that when a nucleic acid sequence and a zinc finger  
protein are aligned according to convention, the primary interaction of the zinc finger is  
with the - strand of the nucleic acid, since it is this strand which is aligned 3' to 5'.  
30 These conventions are followed in the nomenclature used herein. It should be noted,  
however, that in nature certain fingers, such as finger 4 of the protein GLI, bind to the

+ strand of nucleic acid: see Suzuki *et al.*, (1994) NAR 22:3397-3405 and Pavletich and Pabo, (1993) Science 261:1701-1707. The incorporation of such fingers into nucleic acid binding molecules according to the invention is envisaged.

- 5 A zinc finger binding motif is a structure well known to those in the art and defined in, for example, Miller *et al.*, (1985) EMBO J. 4:1609-1614; Berg (1988) PNAS (USA) 85:99-102; Lee *et al.*, (1989) Science 245:635-637; see International patent applications WO 96/06166 and WO 96/32475, corresponding to USSN 08/422,107, incorporated herein by reference.

10

As used herein, "nucleic acid" refers to both RNA and DNA, constructed from natural nucleic acid bases or synthetic bases, or mixtures thereof. Preferably, however, the binding proteins of the invention are DNA binding proteins.

- 15 The structure of the framework of Cys2-His2 zinc fingers is known in the art. The present invention encompasses both those structures which have been observed in nature, including consensus structures derived therefrom, and artificial structures which have non-natural residue numbers and spacing but which retain the functionality of a zinc finger.

20

In general, a preferred zinc finger framework has the structure:




- 25 where X is any amino acid, and the numbers in subscript indicate the possible numbers of residues represented by X.

In a preferred aspect of the present invention, zinc finger nucleic acid binding motifs may be represented as motifs having the following primary structure:

30





wherein X (including  $X^a$ ,  $X^b$  and  $X^c$ ) is any amino acid.  $X_{2-4}$  and  $X_{2-3}$  refer to the presence of 2 or 4, or 2 or 3, amino acids, respectively. The Cys and His residues, which together co-ordinate the zinc metal atom, are marked in bold text and are usually invariant, as is the Leu residue at position +4 in the  $\alpha$ -helix.

Modifications to this representation may occur or be effected without necessarily abolishing zinc finger function, by insertion, mutation or deletion of amino acids. For example it is known that the second His residue may be replaced by Cys (Krizek *et al.*, (1991) J. Am. Chem. Soc. 113:4518-4523) and that Leu at +4 can in some circumstances be replaced with Arg. The Phe residue before  $X_c$  may be replaced by any aromatic other than Trp. Moreover, experiments have shown that departure from the preferred structure and residue assignments for the zinc finger are tolerated and may even prove beneficial in binding to certain nucleic acid sequences. Even taking this into account, however, the general structure involving an  $\alpha$ -helix co-ordinated by a zinc atom which contacts four Cys or His residues, does not alter. As used herein, structures (A) and (B) above are taken as an exemplary structure representing all zinc finger structures of the Cys2-His2 type.

Preferably,  $X^a$  is  $F/Y-X$  or  $P-F/Y-X$ . In this context, X is any amino acid. Preferably, in this context X is E, K, T or S. Less preferred but also envisaged are Q, V, A and P. The remaining amino acids remain possible.

Preferably,  $X_{2-4}$  consists of two amino acids rather than four. The first of these amino acids may be any amino acid, but S, E, K, T, P and R are preferred. Advantageously, it is P or R. The second of these amino acids is preferably E, although any amino acid may be used.

Preferably,  $X^b$  is T or I.

Preferably, X<sup>c</sup> is S or T.

Preferably, X<sub>2-3</sub> is G-K-A, G-K-C, G-K-S or G-K-G. However, departures from the  
5 preferred residues are possible, for example in the form of M-R-N or M-R.

Preferably, the linker is T-G-E-K or T-G-E-K-P.

As set out above, the major binding interactions occur with amino acids -1, +3 and  
10 +6. Amino acids +4 and +7 are largely invariant. The remaining amino acids may  
be essentially any amino acids. Preferably, position +9 is occupied by Arg or Lys.  
Advantageously, positions +1, +5 and +8 are not hydrophobic amino acids, that is to  
say are not Phe, Trp or Tyr. Preferably, position +2 is any amino acid, and preferably  
15 serine, save where its nature is dictated by its role as a ++2 amino acid for an N-  
terminal zinc finger in the same nucleic acid binding molecule.

In a most preferred aspect, therefore, bringing together the above, the invention allows  
the definition of every residue in a zinc finger nucleic acid binding motif which will  
bind specifically to a given nucleic acid triplet.

20

The code provided by the present invention is not entirely rigid; certain choices are  
provided. For example, positions +1, +5 and +8 may have any amino acid  
allocation, whilst other positions may have certain options: for example, the present  
rules provide that, for binding to a central T residue, any one of Ala, Ser or Val may  
25 be used at +3. In its broadest sense, therefore, the present invention provides a very  
large number of proteins which are capable of binding to every defined target nucleic  
acid triplet. As set forth below, these protein may be selected for binding ability using  
polysome display techniques.

30 Preferably, however, the number of possibilities may be significantly reduced before  
selection. For example, the non-critical residues +1, +5 and +8 may be occupied by

the residues Lys, Thr and Gln respectively as a default option. In the case of the other choices, for example, the first-given option may be employed as a default. Thus, the code according to the present invention allows the design of a single, defined polypeptide (a "default" polypeptide) which will bind to its target triplet.

5

In a further aspect of the present invention, there is provided a method for preparing a nucleic acid binding protein of the Cys2-His2 zinc finger class capable of binding to a target nucleic acid sequence, comprising the steps of:

10 a) selecting a model zinc finger domain from the group consisting of naturally occurring zinc fingers and consensus zinc fingers;

b) varying at least one of positions -1, +3, +6 (and ++2) the finger as required according to the rules set forth above; and

15

c) selecting the variants which bind to the target nucleic acid by polysome display.

In general, naturally occurring zinc fingers may be selected from those fingers for which the nucleic acid binding specificity is known. For example, these may be the  
20 fingers for which a crystal structure has been resolved: namely Zif 268 (Elrod-Erickson *et al.*, (1996) Structure 4:1171-1180), GLI (Pavletich and Pabo, (1993) Science 261:1701-1707), Tramtrack (Fairall *et al.*, (1993) Nature 366:483-487) and YY1 (Houbaviy *et al.*, (1996) PNAS (USA) 93:13577-13582).

25 The naturally occurring zinc finger 2 in Zif 268 makes an excellent starting point from which to engineer a zinc finger and is preferred.

Consensus zinc finger structures may be prepared by comparing the sequences of known zinc fingers, irrespective of whether their binding domain is known. Preferably,  
30 the consensus structure is selected from the group consisting of the consensus structure

P Y K C P E C G K S F S Q K S D L V K H Q R T H T G, and the consensus structure P Y K C S E C G K A F S Q K S N L T R H Q R I H T G E K P.

The consensuses are derived from the consensus provided by Krizek *et al.*, (1991) J. Am. Chem. Soc. 113:4518-4523 and from Jacobs, (1993) PhD thesis, University of Cambridge, UK. In both cases, the linker sequences described above for joining two zinc finger motifs together, namely TGEK or TGEKP can be formed on the ends of the consensus. Thus, a P may be removed where necessary, or, in the case of the consensus terminating T G, E K (P) can be added.

10

When the nucleic acid specificity of the model finger selected is known, the mutation of the finger in order to modify its specificity to bind to the target nucleic acid may be directed to residues known to affect binding to bases at which the natural and desired targets differ. Otherwise, mutation of the model fingers should be concentrated upon residues -1, +3, +6 and ++2 as provided for in the foregoing rules.

15

In order to produce a binding protein having improved binding, moreover, the rules provided by the present invention may be supplemented by physical or virtual modelling of the protein/nucleic acid interface in order to assist in residue selection.

20

Zinc finger binding motifs designed according to the invention may be combined into nucleic acid binding proteins having a multiplicity of zinc fingers. Preferably, the proteins have at least two zinc fingers. In nature, zinc finger binding proteins commonly have at least three zinc fingers, although two-zinc finger proteins such as Tramtrack are known. The presence of at least three zinc fingers is preferred. Binding proteins may be constructed by joining the required fingers end to end, N-terminus to C-terminus. Preferably, this is effected by joining together the relevant nucleic acid coding sequences encoding the zinc fingers to produce a composite coding sequence encoding the entire binding protein. The invention therefore provides a method for producing a nucleic acid binding protein as defined above, wherein the nucleic acid

30

binding protein is constructed by recombinant DNA technology, the method comprising the steps of:

- 5 a) preparing a nucleic acid coding sequence encoding two or more zinc finger binding motifs as defined above, placed N-terminus to C-terminus;
- b) inserting the nucleic acid sequence into a suitable expression vector; and
- c) expressing the nucleic acid sequence in a host organism in order to obtain the nucleic acid binding protein.

- 10 A "leader" peptide may be added to the N-terminal finger. Preferably, the leader peptide is MAEEKP.

The nucleic acid encoding the nucleic acid binding protein according to the invention can be incorporated into vectors for further manipulation. As used herein, vector (or  
15 plasmid) refers to discrete elements that are used to introduce heterologous nucleic acid into cells for either expression or replication thereof. Selection and use of such vehicles are well within the skill of the person of ordinary skill in the art. Many vectors are available, and selection of appropriate vector will depend on the intended use of the vector, i.e. whether it is to be used for DNA amplification or for nucleic acid  
20 expression, the size of the DNA to be inserted into the vector, and the host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the host cell for which it is compatible. The vector components generally include, but are not limited to, one or more of the following: an origin of replication, one or more marker genes, an enhancer  
25 element, a promoter, a transcription termination sequence and a signal sequence.

Both expression and cloning vectors generally contain nucleic acid sequence that enable the vector to replicate in one or more selected host cells. Typically in cloning vectors, this sequence is one that enables the vector to replicate independently of the host  
30 chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast and viruses.

The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 $\mu$  plasmid origin is suitable for yeast, and various viral origins (e.g. SV 40, polyoma, adenovirus) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors unless these are used in mammalian cells competent for high level DNA replication, such as COS cells.

Most expression vectors are shuttle vectors, i.e. they are capable of replication in at least one class of organisms but can be transfected into another organism for expression. For example, a vector is cloned in *E. coli* and then the same vector is transfected into yeast or mammalian cells even though it is not capable of replicating independently of the host cell chromosome. DNA may also be replicated by insertion into the host genome. However, the recovery of genomic DNA encoding the nucleic acid binding protein is more complex than that of exogenously replicated vector because restriction enzyme digestion is required to excise nucleic acid binding protein DNA. DNA can be amplified by PCR and be directly transfected into the host cells without any replication component.

Advantageously, an expression and cloning vector may contain a selection gene also referred to as selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that confer resistance to antibiotics and other toxins, e.g. ampicillin, neomycin, methotrexate or tetracycline, complement auxotrophic deficiencies, or supply critical nutrients not available from complex media.

As to a selective gene marker appropriate for yeast, any marker gene can be used which facilitates the selection for transformants due to the phenotypic expression of the marker gene. Suitable markers for yeast are, for example, those conferring resistance to antibiotics G418, hygromycin or bleomycin, or provide for prototrophy in an auxotrophic yeast mutant, for example the URA3, LEU2, LYS2, TRP1, or HIS3 gene.

Since the replication of vectors is conveniently done in *E. coli*, an *E. coli* genetic marker and an *E. coli* origin of replication are advantageously included. These can be obtained from *E. coli* plasmids, such as pBR322, Bluescript® vector or a pUC plasmid, e.g. pUC18 or pUC19, which contain both *E. coli* replication origin and *E. coli* genetic marker conferring resistance to antibiotics, such as ampicillin.

Suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up nucleic acid binding protein nucleic acid, such as dihydrofolate reductase (DHFR, methotrexate resistance), thymidine kinase, or genes conferring resistance to G418 or hygromycin. The mammalian cell transformants are placed under selection pressure which only those transformants which have taken up and are expressing the marker are uniquely adapted to survive. In the case of a DHFR or glutamine synthase (GS) marker, selection pressure can be imposed by culturing the transformants under conditions in which the pressure is progressively increased, thereby leading to amplification (at its chromosomal integration site) of both the selection gene and the linked DNA that encodes the nucleic acid binding protein. Amplification is the process by which genes in greater demand for the production of a protein critical for growth, together with closely associated genes which may encode a desired protein, are reiterated in tandem within the chromosomes of recombinant cells. Increased quantities of desired protein are usually synthesised from thus amplified DNA.

Expression and cloning vectors usually contain a promoter that is recognised by the host organism and is operably linked to nucleic acid binding protein encoding nucleic acid. Such a promoter may be inducible or constitutive. The promoters are operably linked to DNA encoding the nucleic acid binding protein by removing the promoter from the source DNA by restriction enzyme digestion and inserting the isolated promoter sequence into the vector. Both the native nucleic acid binding protein promoter sequence and many heterologous promoters may be used to direct amplification and/or expression of nucleic acid binding protein encoding DNA.

Promoters suitable for use with prokaryotic hosts include, for example, the  $\beta$ -lactamase and lactose promoter systems, alkaline phosphatase, the tryptophan (trp) promoter system and hybrid promoters such as the tac promoter. Their nucleotide sequences have  
5 been published, thereby enabling the skilled worker operably to ligate them to DNA encoding nucleic acid binding protein, using linkers or adapters to supply any required restriction sites. Promoters for use in bacterial systems will also generally contain a Shine-Delgarno sequence operably linked to the DNA encoding the nucleic acid binding protein.

10

Preferred expression vectors are bacterial expression vectors which comprise a promoter of a bacteriophage such as phage  $\lambda$  or T7 which is capable of functioning in the bacteria. In one of the most widely used expression systems, the nucleic acid encoding the fusion protein may be transcribed from the vector by T7 RNA polymerase  
15 (Studier et al, Methods in Enzymol. 185; 60-89, 1990). In the *E. coli* BL21(DE3) host strain, used in conjunction with pET vectors, the T7 RNA polymerase is produced from the  $\lambda$ -lysogen DE3 in the host bacterium, and its expression is under the control of the IPTG inducible lac UV5 promoter. This system has been employed successfully for over-production of many proteins. Alternatively the polymerase gene may be  
20 introduced on a lambda phage by infection with an int- phage such as the CE6 phage which is commercially available (Novagen, Madison, USA). other vectors include vectors containing the lambda PL promoter such as PLEX (Invitrogen, NL) , vectors containing the trc promoters such as pTrcHisXpressTm (Invitrogen) or pTrc99 (Pharmacia Biotech, SE) or vectors containing the tac promoter such as pKK223-3  
25 (Pharmacia Biotech) or PMAL (New England Biolabs, MA, USA).

Moreover, the nucleic acid binding protein gene according to the invention preferably includes a secretion sequence in order to facilitate secretion of the polypeptide from bacterial hosts, such that it will be produced as a soluble native peptide rather than in  
30 an inclusion body. The peptide may be recovered from the bacterial periplasmic space, or the culture medium, as appropriate.



Suitable promoting sequences for use with yeast hosts may be regulated or constitutive and are preferably derived from a highly expressed yeast gene, especially a *Saccharomyces cerevisiae* gene. Thus, the promoter of the TRP1 gene, the ADHI or ADHII gene, the acid phosphatase (PH05) gene, a promoter of the yeast mating pheromone genes coding for the  $\alpha$ - or  $\alpha$ -factor or a promoter derived from a gene encoding a glycolytic enzyme such as the promoter of the enolase, glyceraldehyde-3-phosphate dehydrogenase (GAP), 3-phospho glycerate kinase (PGK), hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triose phosphate isomerase, phosphoglucose isomerase or glucokinase genes, or a promoter from the TATA binding protein (TBP) gene can be used. Furthermore, it is possible to use hybrid promoters comprising upstream activation sequences (UAS) of one yeast gene and downstream promoter elements including a functional TATA box of another yeast gene, for example a hybrid promoter including the UAS(s) of the yeast PH05 gene and downstream promoter elements including a functional TATA box of the yeast GAP gene (PH05-GAP hybrid promoter). A suitable constitutive PH05 promoter is e.g. a shortened acid phosphatase PH05 promoter devoid of the upstream regulatory elements (UAS) such as the PH05 (-173) promoter element starting at nucleotide -173 and ending at nucleotide -9 of the PH05 gene.

Nucleic acid binding protein gene transcription from vectors in mammalian hosts may be controlled by promoters derived from the genomes of viruses such as polyoma virus, adenovirus, fowlpox virus, bovine papilloma virus, avian sarcoma virus, cytomegalovirus (CMV), a retrovirus and Simian Virus 40 (SV40), from heterologous mammalian promoters such as the actin promoter or a very strong promoter, e.g. a ribosomal protein promoter, and from the promoter normally associated with nucleic acid binding protein sequence, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding nucleic acid binding protein by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are relatively orientation and position independent. Many enhancer sequences are known from mammalian genes (e.g. elastase and globin). However, typically one will employ  
5 an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270) and the CMV early promoter enhancer. The enhancer may be spliced into the vector at a position 5' or 3' to nucleic acid binding protein DNA, but is preferably located at a site 5' from the promoter.

10 Advantageously, a eukaryotic expression vector encoding a nucleic acid binding protein according to the invention may comprise a locus control region (LCR). LCRs are capable of directing high-level integration site independent expression of transgenes integrated into host cell chromatin, which is of importance especially where the nucleic acid binding protein gene is to be expressed in the context of a permanently-transfected  
15 eukaryotic cell line in which chromosomal integration of the vector has occurred, or in transgenic animals.

Eukaryotic vectors may also contain sequences necessary for the termination of transcription and for stabilising the mRNA. Such sequences are commonly available  
20 from the 5' and 3' untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding nucleic acid binding protein.

An expression vector includes any vector capable of expressing nucleic acid binding  
25 protein nucleic acids that are operatively linked with regulatory sequences, such as promoter regions, that are capable of expression of such DNAs. Thus, an expression vector refers to a recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector, that upon introduction into an appropriate host cell, results in expression of the cloned DNA. Appropriate expression vectors are well  
30 known to those with ordinary skill in the art and include those that are replicable in eukaryotic and/or prokaryotic cells and those that remain episomal or those which

integrate into the host cell genome. For example, DNAs encoding nucleic acid binding protein may be inserted into a vector suitable for expression of cDNAs in mammalian cells, e.g. a CMV enhancer-based vector such as pEVRF (Matthias, et al., (1989) NAR 17, 6418).

5

Particularly useful for practising the present invention are expression vectors that provide for the transient expression of DNA encoding nucleic acid binding protein in mammalian cells. Transient expression usually involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector, and, in turn, synthesises high levels of nucleic acid binding protein. For the purposes of the present invention, transient expression systems are useful e.g. for identifying nucleic acid binding protein mutants, to identify potential phosphorylation sites, or to characterise functional domains of the protein.

10

Construction of vectors according to the invention employs conventional ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required. If desired, analysis to confirm correct sequences in the constructed plasmids is performed in a known fashion. Suitable methods for constructing expression vectors, preparing in vitro transcripts, introducing DNA into host cells, and performing analyses for assessing nucleic acid binding protein expression and function are known to those skilled in the art. Gene presence, amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA, dot blotting (DNA or RNA analysis), or in situ hybridisation, using an appropriately labelled probe which may be based on a sequence provided herein. Those skilled in the art will readily envisage how these methods may be modified, if desired.

20

25

In accordance with another embodiment of the present invention, there are provided cells containing the above-described nucleic acids. Such host cells such as prokaryote, yeast and higher eukaryote cells may be used for replicating DNA and producing the nucleic acid binding protein. Suitable prokaryotes include eubacteria, such as Gram-

30

negative or Gram-positive organisms, such as *E. coli*, e.g. *E. coli* K-12 strains, DH5a and HB101, or Bacilli. Further hosts suitable for the nucleic acid binding protein encoding vectors include eukaryotic microbes such as filamentous fungi or yeast, e.g. *Saccharomyces cerevisiae*. Higher eukaryotic cells include insect and vertebrate cells, particularly mammalian cells including human cells, or nucleated cells from other multicellular organisms. In recent years propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are epithelial or fibroblastic cell lines such as Chinese hamster ovary (CHO) cells, NIH 3T3 cells, HeLa cells or 293T cells. The host cells referred to in this disclosure  
10 comprise cells in *in vitro* culture as well as cells that are within a host animal.

DNA may be stably incorporated into cells or may be transiently expressed using methods known in the art. Stably transfected mammalian cells may be prepared by transfecting cells with an expression vector having a selectable marker gene, and  
15 growing the transfected cells under conditions selective for cells expressing the marker gene. To prepare transient transfectants, mammalian cells are transfected with a reporter gene to monitor transfection efficiency.

To produce such stably or transiently transfected cells, the cells should be transfected  
20 with a sufficient amount of the nucleic acid binding protein-encoding nucleic acid to form the nucleic acid binding protein. The precise amounts of DNA encoding the nucleic acid binding protein may be empirically determined and optimised for a particular cell and assay.

Host cells are transfected or, preferably, transformed with the above-captioned expression or cloning vectors of this invention and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. Heterologous DNA may be introduced into host cells by any method known in the art, such as transfection with a  
30 vector encoding a heterologous DNA by the calcium phosphate coprecipitation technique or by electroporation. Numerous methods of transfection are known to the

skilled worker in the field. Successful transfection is generally recognised when any indication of the operation of this vector occurs in the host cell. Transformation is achieved using standard techniques appropriate to the particular host cells used.

- 5 Incorporation of cloned DNA into a suitable expression vector, transfection of eukaryotic cells with a plasmid vector or a combination of plasmid vectors, each encoding one or more distinct genes or with linear DNA, and selection of transfected cells are well known in the art (see, e.g. Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press).

10

Transfected or transformed cells are cultured using media and culturing methods known in the art, preferably under conditions, whereby the nucleic acid binding protein encoded by the DNA is expressed. The composition of suitable media is known to those in the art, so that they can be readily prepared. Suitable culturing media are also

15 commercially available.

The zinc finger polypeptides according to the present invention are varied at at least one position. Preferably, the positions selected for variation is one of the positions identified above as being important in determining the binding specificity of the zinc

20 finger polypeptide of the invention.

By "vary" (including grammatical modifications) it is intended to denote that a particular amino acid in the molecule is replaced with an amino acid selected from a varied group, to produce a repertoire of homologous zinc finger polypeptides which

25 differ at the particular amino acid position. The variant amino acids may be selected from a small group of two or more amino acids, from a larger group or may be completely randomly selected from all 20 naturally occurring amino acids. In a preferred embodiment, amino acid analogues and artificial amino acids may be employed.

30

Variation of the zinc finger binding motifs produced according to the invention is preferably directed to those residues where the code provided herein gives a choice of residues. For example, therefore, positions +1, +5 and +8 are advantageously randomised, whilst preferably avoiding hydrophobic amino acids; positions involved in  
5 binding to the nucleic acid, notably -1, +2, +3 and +6, may be randomised also, preferably within the choices provided by the rules of the present invention.

Preferably, therefore, the "default" protein produced according to the rules provided by the invention can be improved by subjecting the protein to one or more rounds of  
10 variation and selection within the specified parameters.


Mutagenesis of zinc finger polypeptides may be achieved by any suitable means. Preferably, the mutagenesis is performed at the nucleic acid level, for example by synthesising novel genes encoding mutant proteins and expressing these to obtain a  
15 variety of different proteins. Alternatively, existing genes can be themselves mutated, such by site-directed or random mutagenesis, in order to obtain the desired mutant genes.

Mutations may be performed by any method known to those of skill in the art.  
20 Preferred, however, is site-directed mutagenesis of a nucleic acid sequence encoding the protein of interest. A number of methods for site-directed mutagenesis are known in the art, from methods employing single-stranded phage such as M13 to PCR-based techniques (see "PCR Protocols: A guide to methods and applications", M.A. Innis, D.H. Gelfand, J.J. Sninsky, T.J. White (eds.). Academic Press, New York, 1990).  
25 Preferably, the commercially available Altered Site II Mutagenesis System (Promega) may be employed, according to the directions given by the manufacturer.

Selection of varied polypeptides according to the invention is carried out by polysome display (see Table 1). This technique relies on coupled transcription and translation of  
30 the coding sequences encoding the zinc finger polypeptides of the invention. This is achieved by preventing dissociation of the mRNA template and the polypeptide chain

from the ribosome, such that the whole entity can be isolated as a polysome. Polysomes are then selected by binding the polypeptide to target nucleic acid, and mRNA eluted from those polysomes which display the desired binding characteristics.

Table 1



The Polysome Display Procedure



Polysome display may be performed according to the methods known in the art, as described below. For example, reference is made to WO95/11922, the methods of which are incorporated herein by reference. The methods of WO95/11922 may be adapted to the present invention, as follows:

5

*Improved Methods For Screening Nascent Peptide Libraries*

A polysome library displaying nascent zinc finger polypeptides can be generated by a variety of methods. Generally, an in vitro translation system is employed to generate polysomes from a population of added mRNA species. Often, the in vitro translation system used is a conventional eukaryotic translation system (e.g., rabbit reticulocyte lysate, wheat germ extract). However, an *E. coli* S30 system (Promega, Madison, Wisconsin) can be used to generate the polysome library from a population of added mRNA species or by coupled transcription/translation (infra). Suitable *E. coli* S30 systems may be produced by conventional methods or may be obtained from commercial sources (Promega, Madison, Wisconsin). The *E. coli* S30 translation system is generally more efficient at producing polysomes suitable for affinity screening of displayed nascent peptides, and the like. Moreover, a prokaryotic translation system, such as the *E. coli* S30 system, has the further advantage that a variety of drugs which block prokaryotic translation (e.g., inhibitors of ribosome function), such as rifampicin or chloramphenicol, can be added at a suitable concentration and/or timepoint to stall translation and produce a population of stalled polysomes, suitable for affinity screening against a target nucleic acid sequence.

In general, the method comprises the steps of: (1) introducing a population of mRNA species into a prokaryotic in vitro translation system (e.g., *E. coli* S30) under conditions suitable for translation to form a pool of polysomes displaying nascent zinc finger polypeptides, so-called polysome forming conditions; (2) contacting the polysomes with a target nucleic acid under suitable binding conditions (i.e., for specific binding to the target nucleic acid and for preserving intact polysome structure); (3) selecting polysomes which are specifically bound to the nucleic acid (e.g., by removing unbound polysomes by washing with a solution); and (4) determining the polynucleotide

30

sequences of the selected polysomes (e.g., by synthesizing cDNA or reverse transcriptase PCR amplification product, and sequencing said cDNA or amplification product). Often, the nucleic acid used for screening is immobilized, such as by being bound to a solid support.

5

In a variation of the method, the population of mRNA molecules is introduced into the in vitro translation system by de novo synthesis of the mRNA from a DNA template. In this improvement, a population of DNA templates capable of being transcribed in vitro (e.g., having an operably linked T7 or SP6 or other suitable promoter) are introduced into a coupled in vitro transcription/translation system (e.g., an *E. coli* S30 system) under conditions suitable for in vitro transcription and translation of the transcribed product. Generally, using a coupled in vitro transcription/translation system is highly efficient for producing polysomes displaying nascent zinc finger polypeptides suitable for affinity screening. Of course, and as noted above, uncoupled systems may also be used, i.e., by adding mRNA to an in vitro translation extract.

10  
15

A further improvement to the general methods of screening nascent zinc finger polypeptide-displaying polysomes comprises the additional step of adding a preblocking agent (e.g., nonfat milk, serum albumin, tRNA, and/or gelatin) prior to or concomitant with the step of contacting the nascent peptide-displaying polysomes with an immobilized nucleic acid. The additional step of adding a preblocking agent reduces the amount of polysomes which bind nonspecifically to the target nucleic acid and/or to the immobilization surface (e.g., microtitre well), thereby enhancing the specificity of selection for polysomes displaying peptides that specifically bind to the nucleic acid. Although the preblocking agent can be selected from a broad group of suitable compositions, the group of preblocking agents comprising: nonfat milk/nonfat milk solids, casein, bovine serum albumin, transfer RNA, and gelatin are preferred, with nonfat milk being especially preferable. Other suitable preblocking agents can be used.

20  
25

Preblocking agents that do not substantially interfere with specific binding (i.e., non-interfering) are suitable.

30

A further improvement to the general methods of screening nascent peptide-displaying polysomes comprises the additional step of isolating polysomes from an in vitro translation reaction (or a coupled in vitro transcription/translation reaction) prior to the step of contacting the nascent peptide-displaying polysomes with nucleic acid. Generally, the polysomes are isolated from a translation reaction by high speed centrifugation to pellet the polysomes, so that the polysome pellet is recovered and the supernatant containing contaminants is discarded. The polysome pellet is resolubilised in a suitable solution to retain intact polysomes. The resolubilised polysomes may be recentrifuged at lower speed (i.e., which does not pellet polysomes) so that the insoluble contaminants pellet and are discarded and the supernatant containing soluble polysomes is recovered, and the supernatant used for affinity screening. Alternatively, the resolubilised polysomes may be used for affinity screening directly (i.e., without low speed centrifugation). Furthermore, the order of centrifugation may be reversed, so that low speed centrifugation is performed prior to high speed centrifugation; the low speed centrifugation supernatant is then centrifuged at high speed and the pelleted polysomes are resolubilised and used for affinity screening. Multiple rounds of high speed and/or low speed centrifugation may be used to increasingly purify the polysomes prior to contacting the polysomes with the immobilized nucleic acid.

Another improvement to the general methods of affinity screening of nascent peptide-displaying polysomes comprises adding a non-ionic detergent to the binding and/or wash buffers. Non-ionic detergent (e.g., Triton X-100, NP-40, Tween, etc.) is added in the binding buffer (i.e., the aqueous solution present during the step of contacting the polysomes with the immobilized nucleic acid) and/or the wash buffer (i.e., the aqueous solution used to wash the bound polysomes (i.e., bound to the immobilized nucleic acid)). Generally, the non-ionic detergent is added to a final concentration of about between 0.01 to 0.5% (v/v), with 0.1% being typical.

Another improvement to the general methods of affinity screening of nascent peptide libraries is generating the DNA template library (from which the mRNA population is

transcribed) in vitro without cloning the library in host cells. Cloning libraries in host cells frequently diminishes the diversity of the library and may skew the distribution of the relative abundance of library members. In vitro library construction generally comprises ligating each member of a population of polynucleotides encoding library members to a polynucleotide sequence comprising a promoter suitable for in vitro transcription (e.g., T7 promoter and leader). The resultant population of DNA templates may optionally be purified by gel electrophoresis. The population of DNA templates is then transcribed and translated in vitro, such as by a coupled transcription/translation system (e.g., E. coli S30).

10

A further improvement to the general methods of affinity screening comprises the added step of combining affinity screening of a nascent peptide-displaying polysome, library with screening of a bacteriophage peptide display library (or other, i.e., peptides on plasmids, expression as secreted soluble antibody in host cells, in vitro expression). In this improvement, polysomes are isolated by affinity screening of a nascent peptide-display library. The isolated polysomes are dissociated, and cDNA is made from the mRNA sequences that encoded nascent peptides that specifically bound to the target nucleic acid). The cDNA sequences encoding the nascent peptide binding regions (i.e., the portions which formed binding contacts to the nucleic acid(s); variable segment sequences) are cloned into a suitable bacteriophage peptide display vector (e.g., pAFF6 or other suitable vector). The resultant bacteriophage vectors are introduced into a host cell to produce a library of bacteriophage particles. Each of the phage clones express on their virion surface the polysome derived peptide sequences as fusions to a coat protein (e.g., as an N-terminal fusion to the PIII coat protein). By incorporating the in vitro-enriched peptide sequences from the polysome screening into a bacteriophage display system, it is possible to continue affinity selection for additional rounds. It is also advantageous, because the resultant bacteriophage display libraries can be screened and tested under conditions that might not have been appropriate for the intact polysomes.

30

Another improvement to the methods of affinity screening is the control of display valency (i.e., the average number of functional zinc finger polypeptides displayed per polysome, and the capacity to vary display valency in different rounds of affinity screening. Typically, a high display valency permits many binding contacts between the polysome and nucleic acid, thus affording stable binding for polysomes which encode zinc finger polypeptide species which have relatively weak binding. Hence, a high display valency system allows screening to identify a broader diversity range of zinc finger polypeptides, since even lower affinity zinc finger polypeptides can be selected. Frequently, such low-to-medium affinity zinc finger polypeptides can be superior candidates for generating very high affinity zinc finger polypeptides, by selecting high affinity zinc finger polypeptides from a pool of mutagenised low-to-medium affinity zinc finger polypeptide clones. Thus, affinity sharpening by mutagenesis and subsequent rounds of affinity selection can be used in conjunction with a broader pool of initially selected zinc finger polypeptide sequences if a high display valency method is used. Alternate rounds of high display valency screening and low display valency screening can be performed, in any order, starting from either a high or low valency system, for as many affinity screening rounds as desired, with intervening variation and sequence diversity broadening, if desired. Alternate rounds of affinity screening, wherein a first round consists of screening a zinc finger polypeptide library expressed in a high valency display system, selecting zinc finger polypeptide clones which bind the target nucleic acid, optionally conducting a mutagenesis step to expand the sequence variability of the selected zinc finger polypeptides, expressing the selected zinc finger polypeptide clones in a lower valency display system, and selecting clones which bind the target nucleic acid, can be performed, including various permutations and combinations of multiple screening cycles, wherein each cycle can be of a similar or different display valency. This improvement affords an overall screening program that employs systems which are compatible with switchable valency (i.e., one screening cycle can have a different display valency than the other(s), and can alternate in order).

Display valency can be controlled by a variety of methods, including but not limited to: controlling the average number of nascent peptides per polysome in a polysome-display

system. This can be controlled by any suitable method, including: (1) altering the length of the encoding mRNA sequence to reduce or increase the frequency of translation termination (a longer mRNA will typically display more nascent peptides per polysome than a shorter mRNA encoding sequence), (2) incorporating stalling (i.e.,  
5 infrequently used) codons in the encoding mRNA, typically distal (downstream of) of the zinc finger polypeptide-encoding portion(s), (3) incorporating RNA secondary structure-forming sequences (e.g., hairpin, cruciform, etc.) distal to the zinc finger polypeptide-encoding portion and proximal to (upstream to) the translation termination site, if any, and/or (4) including an antisense polynucleotide (e.g., DNA, RNA,  
10 polyamide nucleic acid) that hybridizes to the mRNA distal to the zinc finger polypeptide-encoding portion and proximal to (and possibly spanning) the translation termination site, if any. The length of the mRNA may be increased to increase display valency, such as by adding additional reading frame sequences downstream of the zinc finger polypeptide-encoding sequences; such additional reading frame sequences can,  
15 for example, encode the sequence  $(-AAVP-)_n$ , where  $n$  is typically at least 1, frequently at least 5 to 10, often at least 15 to 25, and may be at least 50-100, up to approximately 150 to 500 or more, although infrequently a longer stall sequence can be used. Stalling codons (i.e., codons which are slowly translated relative to other codons in a given translation system) can be determined empirically for any translation system, such as by  
20 measuring translation efficiency of mRNA templates which differ only in the presence or relative abundance of particular codons. For example, a set of clones can be evaluated in the chosen translation system; each species or the set has a stalling polypeptide sequence of 25 amino acids, but each stalling polypeptide sequence consists of a repeating series of one codon, such that all translatable codons are represented in  
25 the set. When translated under equivalent conditions, the zinc finger polypeptide species which produce polysomes having the highest valency (e.g., as determined by sedimentation rate, buoyancy, electron microscopic examination, and other diagnostic methods) thereby identify stalling codons as the codon(s) in the stalling polypeptide sequence.

In one embodiment, a stalling polypeptide sequence is distal (3' to) the zinc finger polypeptide-encoding sequence, and comprises  $-(\text{Gly-Gly-Gly-Gly-Ser})_4\text{-A-A-V-P-}$ , or repeats thereof.

- 5 Alternatively, or in combination with the noted variations, the valency of the target nucleic acid may be varied to control the average binding affinity of selected library members. The target nucleic acid can be bound to a surface or substrate at varying densities, such as by including a competitor target nucleic acid, by dilution, or by other method known to those in the art. A high density (valency) of target nucleic acid can  
10 be used to enrich for library members which have relatively low affinity, whereas a low density (valency) can preferentially enrich for higher affinity library members.

- Each of the improvements to the methods of affinity screening may be combined with other compatible improvements. For example, an in vitro transcription/translation  
15 system can be used in conjunction with a library of DNA templates synthesized in vitro (i.e. without cloning in a host cell). The resultant polysomes can be purified by one or more rounds of high-speed and/or low-speed centrifugation. The purified polysomes can be contacted with an immobilized nucleic acid that is preblocked (e.g., with nonfat milk), and a non-ionic detergent may also be present to further reduce nonspecific  
20 binding. The selected polysomes may then be used as templates for synthesizing cDNA which is then cloned into a bacteriophage display vector, such that the variable segments of the nascent peptides are now displayed on bacteriophage.

#### *Amplification, Affinity Enrichment, And Screening*

- 25 A basic method is described for synthesizing a nascent peptide-polysome library in vitro, screening and enrichment of the library for species having desired specific binding properties, and recovery of the nucleotide sequences that encode those peptides of sufficient binding affinity for target nucleic acid sufficient for selection by affinity selection.

The library consists of a population of nascent zinc finger polypeptide library members comprising nascent peptides. After selecting those nascent peptide library members that bind to the nucleic acid with high affinity, the selected complexes are disrupted and the mRNA is recovered and amplified to create DNA copies of the message. Typically  
5 each copy comprises an operably linked in vitro transcription promoter (e.g., T7 or SP6 promoter). The DNA copies are transcribed in vitro to produce mRNA, and the process is repeated to enrich for zinc finger polypeptides that bind with sufficient affinity.

- 10 The following general steps are frequently followed in the method: (1) generate a DNA template which is suitable for in vitro synthesis of mRNA, (2) synthesize mRNA in vitro by transcription of the DNA templates in a coupled transcription/translation system, (3) bind the nascent peptide to a preferably immobilised target nucleic acid, (4) recover and amplify nascent peptide library members which bind the target nucleic acid  
15 and produce DNA templates from the selected library members competent for in vitro transcription.

Each generated DNA template preferably contains a promoter (e.g., T7 or SP6) which is active in an in vitro transcription system. A DNA template generally comprises (1) a  
20 promoter which is functional for in vitro transcription and operably linked to (2) a polynucleotide sequence encoding an mRNA period. Said encoded mRNA comprises a polynucleotide sequence which encodes a polypeptide comprising a zinc finger polypeptide, (2) a polynucleotide sequence to which a DNA primer suitable for priming first-strand cDNA synthesis of the mRNA can bind, and (3) a ribosome-binding site and  
25 other elements necessary for in vitro translatability of the mRNA, and optionally, for mRNA stability and translatable secondary structure, if any.

Following translation, polysome complexes are screened for high-affinity nucleic acid-binding using standard procedures and as described herein.



After selecting those nascent peptide/polynucleotide complexes that bind with sufficient affinity, the polysomes are isolated and ribosomes released by the addition of EDTA sufficient to chelate the  $Mg^{+2}$  present in the buffer. Ribosomes are removed by high-speed centrifugation, and the RNA component is released by phenol extraction, or by  
5 changing the ionic strength, temperature or pH of the binding buffer so as to denature the nascent peptide. A cDNA copy of the mRNA is made using reverse transcriptase, and the cDNA copy is amplified by, the polymerase chain reaction (PCR). The amplified cDNA is added to the in vitro transcription system and the process is repeated to enrich for those peptides that bind with high affinity.

10

15

The invention is further described, for the purposes of illustration only, in the following examples.

20

#### Example 1

##### *Preparation Of A Varied Zinc Finger Polypeptide*

Zinc finger polypeptides incorporating variation at selected positions are constructed in accordance with the preceding instructions, or as described in any one of  
25 GB9710805.4, GB9710806.2, GB9710807.0, GB9710808.8, GB9710809.6, GB9710810.4, GB9710811.2, GB9710812.0 or EP95928576.8, which are incorporated herein by reference.

#### Example 2

30 *The Template Construct*


The construct is similar to that of Mattheakis *et al* (1994) Proc Natl Acad Sci USA, 91, 9022-9026, but with some modifications to increase the efficiency of ribosome stalling.

5 *General Structure*

The general structure of a transcription template suitable for selection of zinc finger polypeptides according to the present invention is shown in Table 2.

Table 2

5                    General Structure of Zinc Finger Transcription Unit



15

The unit contains a bacteriophage T7 RNA polymerase promoter, which drives a coding sequence encoding a zinc finger polypeptide. Appended to the coding sequence is a linker/stalling sequence region which comprises a flexible Gly/Ser linker, an easily translatable region and a stalling region which is composed of codons rare in *E. coli*. Rare codons hold up the translation process and effectively stall the ribosome on the template.

25    *Sequence Information*

*T7 Promoter*

This is the standard bacteriophage T7 RNA polymerase promoter,, having the sequence TA ATA CGA CTA ACT ATA GGG AGA

30

*Ribosome Binding Site*

This is the bacteriophage T7, gene 10 ribosome binding site. (This and the T7 promoter give high efficiency initiation of transcription and translation). It has the sequence AAGGAG.

5

*Zinc Finger Gene*

A zinc finger coding sequence as shown in SEQ. ID. No. 1 is used.

*Linker/Stalling Sequence*

10

The first 3/4 of this sequence is virtually the same as that used by Mattheakis *et al* (1994). This is because it is the principles behind the design which are important. and not the sequence itself.

15

First there is a 31 residue serine-glycine repeat. This serves as a flexible linker, when translated, which ensures that the expressed zinc finger construct has spacial separation, and flexibility with respect to the stalled ribosome.

20

Second there is a series of six Ala-Ala-Val-Pro repeats. This is a standard, relatively easily translated sequence and serves to ensure that the ribosome is stalled after (and not before) the entire flexible (Ser-Gly) linker has emerged from the ribosome. This is relatively important since approximately ten amino acids are covered by the ribosome at any one time.

25

Third comes a short stretch of codons which contain a high proportion of rare codons - with respect to *E. coli* usage, which slow the translational process and cause regular pauses.

30

Fourth, there is added towards the end of the "third region" an additional ribosome stalling sequence. This has been discussed in Gu *et al* (1994) Proc Natl Acad Sci USA, 91, 5612-5616 and Lovett & Rogers (1996) Microbiological Reviews, 60, 366-385.

35

The sequence:

M V K T D K  
ATG GTT AAA ACA GAT AAA

- 5 when translated, interacts with the peptidyl transferase site of the *E. coli* ribosome, causing translational pausing. In the presence of chloramphenicol, this paused state becomes a stalled state.

10 The sequence is found at the beginning of the *cat-86* gene in *E. coli* which gives resistance to this antibiotic.

It has been found that this sequence increases the efficiency of ribosome stalling by between 10% and 20%, when compared to the exact sequence used by Mattheakis *et al* (1994).

15

Finally, there is no translational STOP codon, so ribosomes will pause when they reach the end of the RNA transcript, before dissociating.

### Example 3

#### 20 The Procedure

The template used is produced by PCR and so is linear, double-stranded DNA of approximately 670bps.

25 Transcription is carried out in a coupled transcription and translation system for linear DNA templates. (The *E. coli* 530 extract system for linear DNA - Promega.)

At present, transcription/translation reactions are carried out in 50µl volumes, each primed with approximately one pmole template - (400-500ng, up to 10<sup>12</sup> DNA molecules).

30

The extract system does not contain T7 RNA polymerase, so this is supplemented by adding T7 polymerase enzyme, and endogenous *E. coli* RNA polymerase is inhibited by adding rifampicin.

5	Template (0.5pM/ $\mu$ l)	2 $\mu$ l	
	Rifampicin (50 $\mu$ g/nl)	1	inhibits <i>E. coli</i> RNA polymerase
	BZA (100mM)	$\frac{1}{2}$	inhibits proteases
	ZnCl <sub>2</sub> (20mM)	1 $\frac{1}{4}$	500 $\mu$ M final concentration, for zinc finger folding
10	Amino acid mix	5	
	530 Premix	20	
	530 Extract	15	
	RNasIN	$\frac{1}{2}$	inhibits RNases
	T7 RNA polymerase	1 ( $\geq 1000$ U)	
15	H <sub>2</sub> O	3 $\frac{3}{4}$	
		<hr/> 50 $\mu$ l	

- incubate 25°C/30 minutes.
- 20 - add three volumes of ice cold stalling buffer.
- place on ice/15 minutes.

Incubation is carried out at 25°C, to help inhibit proteases and RNases, but can be done at any temperature up to 37°C.

25

The ribosomes are stalled by adding to the *in vitro* synthesis system three volumes of "stalling/polysome buffer".

#### *Stalling/Polysome Buffer*

30

This is made at 1 $\frac{1}{3}$  x concentrate but its 1 x working concentrations are:

	Tris (pH7.4)	20mM
	KCl	50mM
	MgCl <sub>2</sub>	10mM
	DTT	5mM
5	ZnCl <sub>2</sub>	50μM
	Chloramphenicol	18μg/ml

The actual ingredients, as far as ribosome stalling is concerned, are Mg<sup>2+</sup> and Chloramphenicol. High Mg<sup>2+</sup> concentration (i.e. 10-20mM) greatly increases the affinity of ribosomes for mRNA (Holschuh & Gassen, (1982) J Biol Chem, 257, 1987-1992) and chloramphenicol causes ribosomes to stall, particularly strongly when combined with the MVKTDK sequence. Tests show that up to 50% of mRNA is attached to ribosomes after stalling cf 40% when using the construct of Mattheakis *et al*, 1994.

15

To collect "polysomes":

- spin 90,000 rpm/30 minutes/4°C (all steps from here to the RT-PCR are carried out in the cold room).
- resuspend polysome pellet in 200μl of stalling buffer.
- 20 - incubate on rolling mixer for 20 minutes/4°C.
- spin down 13,000 rpm/5 minutes ( to pellet insoluble material).
- spin down 13,000 rpm/5 minutes ( to pellet insoluble material).
- collect supernatant.

#### 25 Example 4

##### *Affinity Selection Of Polysomes*

The target ds DNA binding site is pre-bound to streptavidin coated wells to give approximately 1mM concentration once polysomes are added.

30 To 200μl polysomes suspension, add 1μg poly d(I-C) competitor DNA

- immediately add to binding site coated wells.
- incubate 30 minutes.
- wash 6 times with "washing buffer".

#### 5 *Washing Buffer*

This is 1 x "stalling buffer", with the addition of 0.1% between 20 and twice the concentration of KCl and  $MgCl_2$ , to help remove non-specifically bound proteins.

- wash 2 times with 1 x stalling buffer.
- 10 - add 100 $\mu$ l of elution buffer.
- incubate 30 minutes with gentle and occasional agitation.

#### *Elution Buffer*

- 15 This is the same as stalling buffer, but without chloramphenicol or  $MgCl_2$ , and with the addition of 20mM EDTA. The EDTA chelates  $Mg^{2+}$  ions and dissociates ribosomes.

#### *Removal Of DNA Contamination*

- 20 To ensure that the next round of selection is not contaminated by template from the previous round, all ds DNA is removed by incubation with DNaseI.

To 100 $\mu$ l eluted mRNA:

- add 4 $\mu$ l 1M  $MgCl_2$  (since DNaseI requires  $Mg^{2+}$ )
- add 2U DNaseI
- 25 - incubate 37°C/15 minutes
- phenol extract
- ethanol precipitate mRNA
- resuspend in 20 $\mu$ l  $H_2O$ .



*Reverse Transcription*

Reverse transcription is then used to create a ss DNA template from the mRNA collected.

5 *PCR*

The ss DNA is then amplified by PCR to give a double-stranded, full-length template which can be used in the next round of selection experiments.

Claims

1. A method for producing a zinc finger nucleic acid binding protein comprising preparing a zinc finger protein according design rules, varying the protein at one or more positions, and selecting variants which bind to a target nucleic acid sequence by polysome display.
2. A method according to claim 1, wherein the protein is varied at one or more positions selected from the group consisting of +1, +5, +8, -1, +2, +3 and +6.
3. A method for producing a zinc finger nucleic acid binding protein comprising an at least partially varied sequence and selecting variants thereof which bind to a target DNA strand, comprising the steps of:
  - (i) preparing a nucleic acid binding protein of the Cys2-His2 zinc finger class capable of binding to a nucleic acid triplet in a target nucleic acid sequence, wherein binding to each base of the triplet by an  $\alpha$ -helical zinc finger nucleic acid binding motif in the protein is determined as follows:
    - a) if the 5' base in the triplet is G, then position +6 in the  $\alpha$ -helix is Arg; or position +6 is Ser or Thr and position ++2 is Asp;
    - b) if the 5' base in the triplet is A, then position +6 in the  $\alpha$ -helix is Gln and ++2 is not Asp;
    - c) if the 5' base in the triplet is T, then position +6 in the  $\alpha$ -helix is Ser or Thr and position ++2 is Asp;
    - d) if the 5' base in the triplet is C, then position +6 in the  $\alpha$ -helix may be any amino acid, provided that position ++2 in the  $\alpha$ -helix is not Asp;
    - e) if the central base in the triplet is G, then position +3 in the  $\alpha$ -helix is His;
    - f) if the central base in the triplet is A, then position +3 in the  $\alpha$ -helix is Asn;
    - g) if the central base in the triplet is T, then position +3 in the  $\alpha$ -helix is Ala, Ser or Val; provided that if it is Ala, then one of the residues at -1 or +6 is a small residue;

- h) if the central base in the triplet is C, then position +3 in the  $\alpha$ -helix is Ser, Asp, Glu, Leu, Thr or Val;
- i) if the 3' base in the triplet is G, then position -1 in the  $\alpha$ -helix is Arg;
- j) if the 3' base in the triplet is A, then position -1 in the  $\alpha$ -helix is Gln;
- 5 k) if the 3' base in the triplet is T, then position -1 in the  $\alpha$ -helix is Asn or Gln;
- l) if the 3' base in the triplet is C, then position -1 in the  $\alpha$ -helix is Asp;
- (ii) varying the resultant polypeptide at at least one position; and
- 10 (iii) selecting the variants which bind to a target nucleic acid sequence by polysome display.
4. A method according to any preceding claim, wherein the or each zinc finger has the general primary structure
- 15
- (A)  $X^a$  C  $X_{2-4}$  C  $X_{2-3}$  F  $X^c$  X X X X L X X H X X  $X^b$  H - linker
- 1 1 2 3 4 5 6 7 8 9
- wherein X (including  $X^a$ ,  $X^b$  and  $X^c$ ) is any amino acid.
- 20
5. A method according to claim 5 wherein  $X^a$  is  $F/Y-X$  or  $P-F/Y-X$ .
6. A method according to claim 4 or claim 5 wherein  $X_{2-4}$  is selected from any one of: S-X, E-X, K-X, T-X, P-X and R-X.
- 25
7. A method according to any one of claims 4 to 6 wherein  $X^b$  is T or I.
8. A method according to any one of claims 4 to 7 wherein  $X_{2-3}$  is G-K-A, G-K-C, G-K-S, G-K-G, M-R-N or M-R.
- 30
9. A method according to any one of claims 4 to 8 wherein the linker is T-G-E-K or T-G-E-K-P.

10. A method according to any one of claims 4 to 9 wherein position +9 is R or K.
11. A method according to any one of claims 4 to 10 wherein positions +1, +5 and +8  
5 are not occupied by any one of the hydrophobic amino acids, F, W or Y.
12. A method according to claim 11 wherein positions +1, +5 and +8 are occupied by the residues K, T and Q respectively.
- 10 13. A method according to any preceding claim, wherein the polysome display technique comprises the steps of:
- (a) introducing a population of mRNA species into an in vitro translation system under conditions suitable for translation to form a pool of polysomes displaying nascent  
15 zinc finger polypeptides;
- (b) contacting the polysomes with a target nucleic acid under suitable binding conditions;
- (c) selecting polysomes which are specifically bound to the nucleic acid; and
- (d) reverse transcribing and amplifying the isolated mRNA.

## SEQUENCE LISTING

&lt;110&gt; Gendaq Limited

&lt;120&gt; Screening System

&lt;130&gt; p3755

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 2

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 264

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:Synthetic DNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(264)

&lt;400&gt; 1

gca gaa gag aag cct ttt cag tgt cga atc tgc atg cgt aac ttc agc 48  
 Ala Glu Glu Lys Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser  
 1 5 10 15

gat cgt agt agt ctt acc cgc cac acg agg acc cac aca ggc gag aag 96  
 Asp Arg Ser Ser Leu Thr Arg His Thr Arg Thr His Thr Gly Glu Lys  
 20 25 30

cct ttt cag tgt cga atc tgc atg cgt aac ttc agc agg agc gat aac 144  
 Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser Arg Ser Asp Asn  
 35 40 45

ctt acg aga cac cta agg acc cac aca ggc gag aag cct ttt cag tgt 192  
 Leu Thr Arg His Leu Arg Thr His Thr Gly Glu Lys Pro Phe Gln Cys  
 50 55 60

cga atc tgc atg cgt aac ttc agg caa gct gat cat ctt caa gag cac 240  
 Arg Ile Cys Met Arg Asn Phe Arg Gln Ala Asp His Leu Gln Glu His  
 65 70 75 80

cta aag acc cac aca ggc gag aag  
 Leu Lys Thr His Thr Gly Glu Lys  
 85

264

&lt;210&gt; 2

&lt;211&gt; 88

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;223&gt; Description of Artificial Sequence: Synthetic DNA

&lt;400&gt; 2

Ala Glu Glu Lys Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser  
 1 5 10 15

Asp Arg Ser Ser Leu Thr Arg His Thr Arg Thr His Thr Gly Glu Lys  
 20 25 30

Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser Arg Ser Asp Asn  
 35 40 45

Leu Thr Arg His Leu Arg Thr His Thr Gly Glu Lys Pro Phe Gln Cys  
 50 55 60

Arg Ile Cys Met Arg Asn Phe Arg Gln Ala Asp His Leu Gln Glu His  
 65 70 75 80

Leu Lys Thr His Thr Gly Glu Lys  
 85

## INTERNATIONAL SEARCH REPORT

Internal Application No.

**PCT/GB 99/03730**

**A. CLASSIFICATION OF SUBJECT MATTER**  
 FD-302 007K14 / 47 C12B

IPC 7 C07K14/47 C12P21/02

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

### C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 06166 A (MEDICAL RES COUNCIL ;CHOO YEN (SG); KLUG AARON (GB); GARCIA ISIDRO) 29 February 1996 (1996-02-29) cited in the application page 11, paragraph 5	1-13
Y	WO 96 32475 A (UNIV WASHINGTON ;CHENG CHENG (CA)) 17 October 1996 (1996-10-17) cited in the application the whole document	1-13
Y	WO 95 19431 A (SCRIPPS RESEARCH INST ;BARBAS CARLOS F III (US); GOTTESFELD JOEL M) 20 July 1995 (1995-07-20) the whole document	1-13

-/-

**X** Further documents are listed in the continuation of box C.

**X** Patent family members are listed in annex.

• **Special categories of cited documents :**

"A" document defining the general state of the art which is not considered to be of particular relevance

**"E"** earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

\* document published prior to the international filing date but later than the priority date claimed

\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

**"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone**

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

**"a" document member of the same patent family**

Date of the actual completion of the international search

**10 March 2000**

Date of mailing of the international search report

**17/03/2000**

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patenslaan 2  
NL - 2260 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

**Authorized officer**

Hillenbrand, G

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 99/03730

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 11922 A (AFFYMAX TECH NV ;DOWER WILLIAM J (US); MATTHEAKIS LARRY C (US)) 4 May 1995 (1995-05-04) cited in the application the whole document	1-13
P,X	WO 98 53059 A (MEDICAL RES COUNCIL ;ISALAN MARK (GB); CHOO YEN (GB); KLUG AARON ( )) 26 November 1998 (1998-11-26) the whole document	1-13
P,X	WO 99 47656 A (MEDICAL RES COUNCIL ;ISALAN MARK (GB); CHOO YEN (GB)) 23 September 1999 (1999-09-23) the whole document	1-13
A	DESJARLAIS J R ET AL: "TOWARD RULES RELATING ZINC FINGER PROTEIN SEQUENCES AND DNA BINDINGSITE PREFERENCES" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA,US,NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 89, no. 16, 1 August 1992 (1992-08-01), pages 7345-7349, XP002009496 ISSN: 0027-8424 cited in the application the whole document	1
A	NARDELLI J ET AL: "ZINC FINGER-DNA RECOGNITION: ANALYSIS OF BASE SPECIFICITY BY SITE-DIRECTED MUTAGENESIS" NUCLEIC ACIDS RESEARCH,GB,OXFORD UNIVERSITY PRESS, SURREY, vol. 20, no. 16, 25 August 1992 (1992-08-25), pages 4137-4144, XP002009497 ISSN: 0305-1048 cited in the application the whole document	1



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Initial Application No

PCT/GB 99/03730

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9606166 A	29-02-1996	AU 1003799 A	22-04-1999
		AU 698152 B	22-10-1998
		AU 3229195 A	14-03-1996
		CA 2196419 A	29-02-1996
		EP 0781331 A	02-07-1997
		JP 10504461 T	06-05-1998
		US 6013453 A	11-01-2000
		US 6007988 A	28-12-1999
WO 9632475 A	17-10-1996	AU 5445296 A	30-10-1996
WO 9519431 A	20-07-1995	AU 704601 B	29-04-1999
		AU 1686595 A	01-08-1995
		CA 2181548 A	20-07-1995
		EP 0770129 A	02-05-1997
		FI 962879 A	17-09-1996
		JP 9508019 T	19-08-1997
		NO 962991 A	18-09-1996
WO 9511922 A	04-05-1995	AU 8124694 A	22-05-1995
		US 5922545 A	13-07-1999
WO 9853059 A	26-11-1998	AU 7542498 A	11-12-1998
WO 9947656 A	23-09-1999	AU 2944999 A	11-10-1999

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**